

Course Introduction & Convex Optimization

R. Kyng & M. Probst

Problem Set 2 — Monday, February 26th

These exercises will not count toward your grade, but you are encouraged to solve them all. This exercise sheet contains exercises relating to lectures in Week 2.

To get feedback, you must hand in your solutions by 23.59 pm on March 7th. Both hand-written and L^AT_EX solutions are acceptable, but we will only attempt to read legible text.

Exercise 1

Prove that if a matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ is symmetric, then $\|\mathbf{A}\| = \max(|\lambda_{\max}(\mathbf{A})|, |\lambda_{\min}(\mathbf{A})|)$ and give an example of a non-symmetric matrix for which this is not true.

Exercise 2

Consider a twice continuously differentiable function $f : S \rightarrow \mathbb{R}$, where $S \subset \mathbb{R}^n$ is a convex open set. Prove that f is β -gradient Lipschitz if and only if for all $\mathbf{x} \in S$ we have $\|\mathbf{H}_f(\mathbf{x})\| \leq \beta$.

Exercise 3

Prove that when running Gradient Descent, $\|\mathbf{x}_i - \mathbf{x}^*\|_2 \leq \|\mathbf{x}_0 - \mathbf{x}^*\|_2$ for all i .

Exercise 4

Prove the following theorem.

Theorem. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be an β -gradient Lipschitz, convex function. Let \mathbf{x}_0 be a given starting point, and let $\mathbf{x}^* \in \arg \min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x})$ be a minimizer of f . The Gradient Descent algorithm given by $\mathbf{x}_{i+1} = \mathbf{x}_i - \frac{1}{\beta} \nabla f(\mathbf{x}_i)$ ensures that the k th iterate satisfies

$$f(\mathbf{x}_k) - f(\mathbf{x}^*) \leq \frac{2\beta \|\mathbf{x}_0 - \mathbf{x}^*\|_2^2}{k+1}.$$

Hint: do an induction on $1/\text{gap}_i$.

Exercise 5

1. For each of the following functions answer these questions:

- Is the function convex?

- Is the function β -gradient Lipschitz for some β ?
- If the function is β -gradient Lipschitz give an upper bound on β – the bound should be within a factor 4 of the true value.

- (a) $f(x) = |x|^{1.5}$ on $x \in \mathbb{R}$
 (b) $f(x) = \exp(x)$ on $x \in \mathbb{R}$
 (c) $f(x) = \exp(x)$ on $x \in (-1, 1)$
 (d) $f(x, y) = \sqrt{x + y}$ on $(x, y) \in (0, 1) \times (0, 1)$.
 (e) $f(x, y) = \sqrt{x + y}$ on $(x, y) \in (1/2, 1) \times (1/2, 1)$.
 (f) $f(x, y) = \sqrt{x^2 + y^2}$ on $(x, y) \in \mathbb{R}^2$.

Bonus Exercise 6: Strongly Convex Functions

This longer exercise will teach you about *strongly convex* functions. In it, you will show that, with a small tweak, gradient descent quickly converges to a highly accurate solutions on these functions.

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function. Assume f is twice continuously (Fréchet) differentiable and that its first and second (Fréchet) derivatives are integrable (basically, don't worry that weird stuff is happening with the derivatives). Assume that for all \mathbf{x} , we have for some constant $\mu > 0$, that $\lambda_{\min}(H_f(\mathbf{x})) \geq \mu$. When this holds, we say that f is μ -strongly convex.

Part A. Prove that for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{\mu}{2} \|\mathbf{y} - \mathbf{x}\|_2^2.$$

Part B. Prove that there is value $L \in \mathbb{R}$ such that for all $\mathbf{x} \in \mathbb{R}^n$, we have $f(\mathbf{x}) \geq L$. In other words, the function is not unbounded below.

Part C. Prove that f is *strictly convex* as per Definition 3.2.8 in Chapter 3. Prove also that the minimizer $\mathbf{x}^* \in \arg \min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x})$ of f is unique.

Part D. Let \mathbf{x}_0 be a given starting point and \mathbf{x}^* be the minimizer of f . Suppose we have an algorithm DECENTDESCENT which takes a starting point \mathbf{x}_0 , and a step count $t \in \mathbb{N}$. DECENTDESCENT(\mathbf{x}_0, t) runs for t steps and returns $\tilde{\mathbf{x}} \in \mathbb{R}^n$ such that

$$f(\tilde{\mathbf{x}}) - f(\mathbf{x}^*) \leq \frac{\gamma \|\mathbf{x}_0 - \mathbf{x}^*\|_2^2}{t + 1}$$

where $\gamma > 0$ is a positive number.

Assume that the *cost* of running DECENTDESCENT for t steps is t . Explain how, with a total cost of at most $\frac{8\gamma}{\mu} \log(\|\mathbf{x}_0 - \mathbf{x}^*\|_2 / \delta)$, we can produce a point $\hat{\mathbf{x}} \in \mathbb{R}^n$ such that $\|\hat{\mathbf{x}} - \mathbf{x}^*\|_2 \leq \delta$ for $\delta > 0$.

Part E. Consider a function $h : \mathbb{R}^n \rightarrow \mathbb{R}$ which is both μ -strongly convex and β -gradient Lipschitz. Give an algorithm that returns \mathbf{x}' with

$$h(\mathbf{x}') - h(\mathbf{x}^*) \leq \epsilon$$

by computing the gradient of h at at most $\frac{32\beta}{\mu} \log(2\beta \|\mathbf{x}_0 - \mathbf{x}^*\|_2^2 / \epsilon)$ points.