

**Solution 1**

- (a) The standard BASICMINCUT algorithm can be implemented using an array containing all the degrees of the vertices currently in the graph — for the purpose of efficiently selecting an edge u.a.r. for contraction. The only adaptation needed now is that after each contraction, we scan this array and maintain a global minimal degree ever seen. It is clear that this can be done in linear time per step and thus in  $\mathcal{O}(n^2)$  time in total.
- (b) We know from the lecture that contractions can only increase the size of a minimum cut but never decrease it. Since the edges incident to any one vertex always *form* a cut, each of the numbers that we could report in this algorithm corresponds to some cut in the original graph, which readily implies the claim.
- (c) For  $n \leq 2$ , the claim is empty. Let us now look at  $n > 2$ , let us fix a graph  $G$  of size  $n$  and a cut  $C$  of size  $\mu(G)$ . What is the probability that the event we are looking for occurs? There are two cases. Either,  $G$  contains a vertex of degree less than  $(1 + \alpha)\mu(G)$ . In that case, no matter what the further recursion would yield, we will always return a number at most that degree and thus the probability is 1. Or, all vertices in  $G$  have degree at least  $(1 + \alpha)\mu(G)$ . Then, there are at least  $(1 + \alpha)\mu(G) \cdot \frac{n}{2}$  edges in the graph and the probability that we contract one from  $C$  is thus bounded by  $\frac{2}{(1+\alpha)n}$ . But then with the complement of this probability, the new graph  $G/e$  will still have a cut of size  $\mu(G)$  and by induction, the claim follows.

As for the calculation (which was not required), since  $p_\alpha(2) = 1$ , we just compute (for

$n$  sufficiently large)

$$\begin{aligned}
p_\alpha(n) &\geq \prod_{i=3}^n \left(1 - \frac{2}{(1+\alpha)i}\right) \\
&= \exp\left(\sum_{i=3}^n \ln\left(1 - \frac{2}{(1+\alpha)i}\right)\right) \\
&\stackrel{(1)}{\geq} \exp\left(\sum_{i=3}^n \left(-\frac{2}{(1+\alpha)i} - \left(\frac{2}{(1+\alpha)i}\right)^2\right)\right) \\
&\stackrel{(2)}{\geq} \exp\left(-\frac{2}{1+\alpha} \left(H_n - \frac{3}{2}\right) - \frac{2\pi^2}{6}\right) \\
&\stackrel{(3)}{\geq} \exp\left(-\frac{2}{1+\alpha} \ln n - \frac{2\pi^2}{6}\right) \\
&= \Omega(n^{\frac{2}{1+\alpha}}),
\end{aligned}$$

where (1) uses the inequality  $1 - x \geq e^{-x-x^2}$  for  $x \leq 0.68$  and  $\frac{2}{(1+\alpha)i} < 2/3 < 0.68$  for all  $\alpha > 0$  and  $i \geq 3$ ; where (2) uses that  $\sum_{i=1}^{\infty} \frac{1}{i^2} = \frac{\pi^2}{6}$  and  $\frac{2}{1+\alpha} < 2$  for  $\alpha > 0$ ; and where (3) uses  $H_n - 3/2 < \ln n$  (the difference between  $\ln n$  and  $H_n$  approaches the Euler-Mascheroni constant  $\gamma$ , which is about 0.577).

## Solution 2

Given a graph  $G = (V, E)$ , let  $N$  be the number of minimum cuts in  $G$ . We want to show that  $N \leq \binom{n}{2}$  where  $n := |V|$ .

Let  $C_1, \dots, C_N$  be the minimum cuts in  $G$ . Then, we know that for each  $i \in \{1..N\}$

$$\Pr[C_i \text{ is found by Karger's algorithm BASICMINCUT}(G)] \geq \frac{1}{\binom{n}{2}}.$$

Now, we observe that for each two distinct indices  $i, j \in \{1..N\}$  the events “ $C_i$  is found by BASICMINCUT( $G$ )” and “ $C_j$  is found by BASICMINCUT( $G$ )” are disjoint (i.e. they never happen at the same time). To see this, consider the graph obtained at the termination of BASICMINCUT( $G$ ). It has only two vertices and these vertices (together with the “contraction history”) uniquely determine a partition of the vertex set  $V$ . So we cannot get two different minimum cuts from one execution of the algorithm. Therefore, it follows that

$$\begin{aligned}
&\Pr[\text{a minimum cut is found by BASICMINCUT}(G)] \\
&= \sum_{i=1}^N \Pr[C_i \text{ is found by BASICMINCUT}(G)] \geq \frac{N}{\binom{n}{2}}.
\end{aligned}$$

Since  $\Pr[\text{a minimum cut is found by BASICMINCUT}(G)] \leq 1$  (because it is a probability!), we obtain  $N \leq \binom{n}{2}$ .

### Solution 3

- (a) Observe that there are three (potentially empty) sets of edges  $e = \{u, v\}$  that are important in this scenario:

$$\begin{aligned} E_1 &:= \{e = \{u, v\} \mid u \in A \cap B \text{ and } v \in V \setminus (A \cup B)\} \\ E_2 &:= \{e = \{u, v\} \mid u \in (A \setminus B) \cup (B \setminus A) \text{ and } v \in V \setminus (A \cup B)\} \\ E_3 &:= \{e = \{u, v\} \mid u \in A \cap B \text{ and } v \in (A \setminus B) \cup (B \setminus A)\}. \end{aligned}$$

It is easy to see that  $f(A \cap B) + f(A \cup B) = |E_1| + |E_3| + |E_1| + |E_2|$ , while  $f(A) + f(B) \geq 2|E_1| + |E_2| + |E_3|$ , the last inequality holding because some edges between  $A \setminus B$  and  $B \setminus A$  may appear both in  $f(A)$  and  $f(B)$ .

- (b) Let  $k$  be the size of a minimum cut. Then by (a) we get  $f(A \cap B) + f(A \cup B) \leq f(A) + f(B) = k + k$ , which together with  $f(C) \geq k$  for any set  $C \neq \emptyset, V$ , implies  $f(A \cap B) = f(A \cup B) = k$ .
- (c) Suppose towards a contradiction that  $S \neq S'$ , with  $S, S' \subset V$  and  $s \in S \cap S'$  are such that  $C(S)$  and  $C(S')$  are both minimum cuts and  $|S| = |S'|$  is minimal. Note that as  $t \notin S \cup S'$  because they are both cuts, we must have  $S \cup S' \neq V$ , and so we can in a similar way to part (b) prove that  $S \cap S'$  is a minimum cut with  $|S \cap S'| < |S| = |S'|$ , a contradiction.

### Solution 4

- (a) Since  $G$  is connected, there are at least  $n - 1$  edges. If there are at least  $n$  edges, then  $\Pr[\mu(G) \neq \mu(G/e)] \leq \frac{1}{n}$  since in the worst case the minimum cut is unique and the probability of contracting a given edge is at most the claimed bound.

If there are only  $n - 1$  edges the graph is a tree. Then contracting any edge keeps the graph a tree with 1 fewer vertex which means that  $\Pr[\mu(G) \neq \mu(G/e)] = 0 \leq \frac{1}{n}$ .

- (b) BASICMINCUT will succeed if it never contracts a given edge in the cut of size 1. This happens with probability at least

$$\begin{aligned} &\left(1 - \frac{1}{n}\right) \left(1 - \frac{1}{n-1}\right) \cdots \left(1 - \frac{1}{3}\right) \\ &= \frac{\binom{n-1}{n}}{\binom{n-1}{n}} \cdots \frac{\binom{2}{3}}{\binom{2}{3}} = \frac{2}{n}. \end{aligned}$$

- (c) A contraction of an edge changes the degree of at most 2 vertices. Therefore there is still at least one vertex of degree  $k \Rightarrow$  minimum cut is still of size  $k$ .

(d) If there are 3 or more vertices of degree  $k$  we are done by (c). If there are 0 or 1 vertices of degree  $k$ , then the number of edges is at least  $((n-1)(k+1)+k)/2 = (n(k+1)-1)/2$  so that when we fix a minimum cut of size  $k$ , the probability of contracting one of the edges of the minimum cut is at most

$$\frac{k}{(n(k+1)-1)/2} = \frac{2k}{n(k+1)-1}.$$

It remains to consider the case that there are two vertices of degree  $k$ . If these two vertices are not adjacent, then there are two disjoint minimum cuts which means that  $\Pr[\mu(G) \neq \mu(G/e)] = 0$ . If they are adjacent, then the minimum cut may only change if we contract one of the at most  $k-1$  edges between the two vertices (they can't be connected via  $k$  edges since the graph is connected and  $n \geq 3$ ). Since there are at least  $(2k + (n-2)(k+1))/2$  edges, the failure probability is at most

$$\frac{2(k-1)}{2k + (n-2)(k+1)} \leq \frac{2k-2}{n(k+1)-2} \leq \frac{2k-1}{n(k+1)-1} \leq \frac{2k}{2n(k+1)-1}.$$

The second to last inequality holds because  $a := 2k-1 \leq n(k+1)-1 =: b$  and then

$$\frac{a-1}{b-1} \leq \frac{a}{b} \Leftrightarrow -b \leq -a \Leftrightarrow a \leq b.$$